

NIKHEF-H/94-09

May 1994

subm. to Phys. Lett. B

Experimental momentum spectra of identified hadrons in jets and the predictions from LPHD + MLLA

N.C. Brümmer

NIKHEF-H

P.O. Box 41882

1009 DB Amsterdam NL

e-mail: Nichol.Brummer@nikhef.nl

Abstract

Experimental data on the shape of hadronic momentum spectra are compared with theoretical predictions in the context of calculations in the Modified Leading Log Approximation (MLLA), under the assumption of Local Parton Hadron Duality (LPHD). Considered are experimental measurements at e^+e^- -colliders of ξ_p^* , the position of the maximum in the distribution of $\xi_p = \log(1/x_p)$, where $x_p = p/p_{\text{beam}}$. The parameter ξ_p^* is determined for various hadrons at various centre of mass energies. It is interesting to look at the dependence of ξ_p^* on the hadron type. This is used to study the influence of the hadron type on the cut-off scale Q_0 in the parton shower development. The dependence of ξ_p^* on the centre of mass energy is seen to be described adequately by perturbation theory. The approach is made quantitative by extracting a value of $\alpha_s(m_Z)$ from an overall fit to the scaling behaviour of ξ_p^* .

1 Introduction

During a few years of LEP running, a large amount of information was collected on identified hadron species in jets. The higher centre of mass energy of LEP, compared to past e^+e^- accelerators, also makes it easier to separate the behaviour of hadrons with a high momentum, which are correlated strongly to the primary quark, from those with a low momentum, created mainly during fragmentation and hadronisation.

Clear scaling violations have been observed in the shape of the charged particle x_p -distribution as function of the centre of mass energy. The strong coupling constant has been extracted from these scaling violations, using the behaviour at high momenta: $0.2 \lesssim x_p \lesssim 0.7$. Due to the larger statistical errors this is not possible for individually identified hadron species.

However, low momentum data for specific types of hadrons maybe be used to study the properties of jet-evolution and hadron-formation in the context of the LPHD hypothesis and MLLA calculations of parton spectra [1–4]. The assumption of ‘Local Parton Hadron Duality’ (LPHD) states that a calculated spectrum for ‘partons’ in a ‘parton shower’ can be related to the spectrum of real hadrons by simple normalisation constants. These constants have to be determined by experiment. A second assumption is that the low momentum part of the spectrum is not influenced in a significant way by hadrons that are correlated to the primary quark.

Calculations of the parton spectra in the ‘Modified Leading Log Approximation’ (MLLA) take into account next-to-leading logarithms in a consistent fashion. The physical mechanism relevant to these next-to-leading logarithms is the coherent emission of soft gluons inside a jet, leading to an angular ordering and an effective transverse momentum cutoff for the partons. Parton jets develop through repeated parton splittings, resulting in an increase of the multiplicity at lower momenta. The interplay of coherent emission of gluons and the creation of hadrons causes this spectrum to be cut off at very low momenta. Calculations predict the shape of the ξ_p distribution. The resulting ‘hump-backed’ distribution is nearly gaussian. As an example, the Monte Carlo spectrum of the Λ baryon in Z^0 decays can be seen in figures 1 and 2. In the following ξ_p^* will be determined for various types of hadrons and




Figure 1: The dependence of the Λ spectrum on the species of its ‘parent’ particle in JETSET. Particles produced directly in the fragmentation process have the Lund-model string as ‘parent’. The scales of both x_p and ξ_p are given for comparison.




Figure 2: The dependence of the Λ spectrum on the flavour of the primary quarks in Z^0 decays at LEP, as predicted by JETSET. Notice that the high and low momentum regions are not separated for Λ ’s in s , c and b events.

	$\sqrt{s}$	charged	$\pi^\pm$	π^0	$K^\pm$	K^0	p	Λ	Ξ^-
ARGUS	9.98		2.32 ± 0.022	2.42 ± 0.08	1.64 ± 0.05	1.72 ± 0.04	1.63 ± 0.07	1.44 ± 0.034	1.32 ± 0.11
CLEO	10.49		2.30 ± 0.08			1.68 ± 0.17	1.67 ± 0.10	1.47 ± 0.08	
TASSO	14	2.45 ± 0.05	2.66 ± 0.06		1.91 ± 0.13	1.65 ± 0.13	1.80 ± 0.15		
JADE	14			2.68 ± 0.27					
TASSO	22	2.74 ± 0.06	2.99 ± 0.09		2.41 ± 0.23	2.47 ± 0.8	2.14 ± 0.27	1.75 ± 0.50	
JADE	22.5			2.84 ± 0.30					
HRS	29		> 3.3		2.28 ± 0.42	2.25 ± 0.30		1.96 ± 0.12	
TPC/ 2γ	29		3.00 ± 0.05		2.14 ± 0.05	1.98 ± 0.09	2.13 ± 0.11		
TASSO	34	3.00 ± 0.05	3.17 ± 0.05		2.35 ± 0.20	2.32 ± 0.10	2.24 ± 0.09	2.15 ± 0.15	
JADE	35			3.24 ± 0.05					
TASSO	44		3.37 ± 0.07		< 3.5		< 3.5		
JADE	44			3.52 ± 0.08					
TOPAZ	58	3.42 ± 0.04	3.51 ± 0.07		2.83 ± 0.10		2.64 ± 0.06		
OPAL	91.2	3.603 ± 0.013				2.91 ± 0.04		2.77 ± 0.05	2.57 ± 0.11
DELPHI	91.2					2.62 ± 0.11		2.81 ± 0.04	
L3	91.2	3.71 ± 0.05		4.11 ± 0.18					

Table 1: The values of ξ_p^* for various hadrons as determined from the momentum spectra as measured by e^+e^- experiments at different values of $\sqrt{s}$. For the η meson, the L3 experiment [10] reported a value $\xi_p^* = 2.60 \pm 0.15$.

at various centre of mass energies. Subsequently a comparison is made with the theoretical calculations.

Figure 1 shows that LPHD is not an obvious assumption: in the JETSET Monte Carlo, the spectrum of e.g. a Λ baryon depends on its ‘parent’ particle. The LPHD assumption states that the sum of these spectra is proportional to the spectrum as calculated for a parton shower with a correctly chosen energy cut-off.

Moreover, the JETSET momentum spectrum of the Λ baryon (figure 2) shows a strong dependence on the flavour of the primary quarks. For the heavier quarks, s, c and b, the high momentum contribution correlated to the primary quarks is not separated from the low momentum contribution related to the fragmentation process. This can be expected to cause shifts in ξ_p^* that depend on the jet-flavour.

The spectrum of all charged particles at LEP has been fitted to the MLLA distribution, with free normalisation factors for pions, kaons and protons, depending on the centre of mass energy [5,6]. It was quite surprising [3] that the MLLA functions can also fit the high momentum part in the data for pions. This is probably a coincidence, since the heavier K^0 meson has a spectrum that can not be fitted as nicely at large x_p .


One way to see whether the observed shape of the spectrum is really due to coherent gluon emission is to compare experimental data to predictions of

the JETSET Monte Carlo program, with the coherence (angular ordering) either turned on or turned off. The large dependence on the primary quark flavours and the fact that coherence in JETSET is not really necessary to fit the experimental data has led some to the conclusion [7] that coherence can not be demonstrated on the basis of hadron momentum spectra. Here an attempt will be made to infer more information using the properties of identified hadrons, concentrating on the observable ξ_p^* . This is done for data from LEP and from various other e^+e^- colliders. Subsequently, the available results for ξ_p^* are compared to the LPHD + MLLA approach [1,2].

2 Experimental values of ξ_p^* for identified hadrons

At LEP, analyses of explicitly identified hadrons have been performed for the π^0 [6], the K_S^0 [8,9], the η [10] and the Λ [8,11,12,22]. At lower statistics data is also available for the Ξ^- and Ω^- baryons, as well as the Σ^* and Ξ^* [8,11].

The values of ξ_p^* correspond to low momenta, where the dependence on the primary quark flavours is expected to be small. To first approximation the distribution in ξ_p is gaussian, but a distorted gaussian fits the calculated spectrum more accurately [13,2]. The present statistics of identified hadrons



(a) ξ_p^* versus the hadron mass (b) Q_0 versus the hadron mass (c) Q_0 versus the hadron mass per quark

Figure 3: The maxima ξ_p^* of the distributions in $\xi = -\log x_p$ for different hadrons as measured at LEP (table 1). the value of ξ_p^* is a nearly linear function of $\log Q_0$. The exact dependence for $\Lambda_{\text{eff}} = 150 \text{ MeV}$ leads to figure (b), where the naive expectation would be that Q_0 is a linear function of the hadron mass. Figure (c) gives the dependence of Q_0 on the hadron mass divided by the number of valence quarks.

is not yet sufficient to extract the width and the two distortion parameters. However, since these other parameters depend more strongly on the high momentum data, they can be expected to depend stronger on the event flavour, which could bias the results. This is a reason not to use a distorted gaussian to determine ξ_p^* .

In the following ξ_p^* was determined by fitting the ξ_p distribution in a limited range of ξ_p around the maximum, to a gaussian distribution. A systematic error was estimated by changing the fitted range. The statistical and systematic errors were added in quadrature. The systematic error often dominated.

The results are given in table 1. Use was made of spectra published by CLEO [15], ARGUS [16], TPC [17], HRS [18], TASSO [19], JADE [20], TOPAZ [21], DELPHI [8,22], OPAL [5,9,11] and L3 [6,10]. Most of the recent publications reported values of ξ_p^* . Most experiments at lower centre of mass energies did not report the distribution of ξ_p or x_p , but used the energy to define $x_E = 2E/E_{\text{beam}}$, or they divided by $\beta = v/c$ to obtain the ‘scaling cross section’. Many different normalisations have been used, but fortunately this is unimportant for a determination ξ_p^* . The experiments (ARGUS and CLEO) near the Υ resonances give results for the ‘continuum’, and for the resonances itself, where the contributions from the continuum are subtracted. The values in table 1 refer to data from the continuum above the Υ resonance. In the data from ARGUS for charged pions and protons,

the contributions from decays of K_S^0 and Λ were subtracted.

3 Predictions about ξ_p^*

The LPHD + MLLA calculations [2] of the distribution of ξ_p depend on an effective QCD scale $\Lambda_{\text{eff}} \sim \Lambda_{\text{QCD}}$ and on a transverse momentum cutoff Q_0 in the evolution of the parton cascade. The value of ξ_p^* is calculated to be a nearly linear function of $\log Q_0$.

The calculated dependence on the centre of mass energy is as follows [2]:

$$\xi_p^* = Y \left[\frac{1}{2} + \sqrt{C/Y} - C/Y + \mathcal{O}(Y^{-3/2}) \right] + F(\lambda) \quad (1)$$

where $F(0) = 0$ and¹

$$Y = \log(E_{\text{beam}}/\Lambda_{\text{eff}}), \quad \lambda = \log(Q_0/\Lambda_{\text{eff}}) \quad (2)$$

and Λ_{eff} is the ‘effective’ QCD scale, while only the cutoff scale Q_0 depends on the hadron type. The constant C is calculated to be

$$C = \left(\frac{a}{4N_c} \right)^2 \frac{N_c}{b}, \quad (3)$$

¹This detail [23] is not completely clear in reference [2], where equation (1) is given as a function of $Y - \lambda$, but in the approximation that $\lambda = 0$.

where

$$b = 11N_c/3 - 2n_f/3, \quad (4)$$

$$a = 11N_c/3 + 2n_f/(3N_c^2). \quad (5)$$

$N_c = 3$ is the number of colours and $n_f = 3$ is the active number of quark flavours in the fragmentation process. For $n_f = 3$ one finds $C = 0.2915$.

It is important to note that F does not depend on Y , and that the first part of equation (1) is independent of Q_0 . This predicted behaviour can be checked by comparing spectra of different identified particles. In the available momentum range this leads to a nearly linear dependence of ξ_p^* on Y .

In ref. [2] various graphs show the results of numerical calculations for the dependence of various parameters on E_{beam} and $Q_0 \neq \Lambda_{\text{eff}}$. Figure 4 of ref. [2] shows the dependence of ξ_p^* on Q_0 for $\Lambda_{\text{eff}} = 150 \text{ MeV}$, and at a number of beam energies. From this it is possible to extract F after subtracting the Y dependent part of equation (1). For an interpretation of the experimental values of ξ_p^* it was useful to fit both this function $F(\lambda)$, and its inverse $\lambda(F)$ to polynomials. The result for $F(\lambda)$ is:

$$F(\lambda) = -1.380 \cdot \lambda + 0.086 \cdot \lambda^2 + 0.044 \cdot \lambda^3 \pm 0.06 \quad (6)$$

while the inverse of F is described by:

$$\lambda(F) = -0.749 \cdot F - 0.054 \cdot F^2 - 0.072 \cdot F^3 \pm 0.06 \quad (7)$$

The given accuracy is the maximum deviation from the distribution plotted in ref. [2], in the range $-2 \lesssim F \lesssim 0$ and $0 \lesssim \lambda \lesssim 2$.

4 Dependence of ξ_p^* on the hadron mass

The dependence of ξ_p^* on the hadron mass as determined from LEP data is shown in figure 3(a). Using equation (7) we can convert the values of ξ_p^* for a given value of $\Lambda_{\text{eff}} = 150 \text{ MeV}$ to values of Q_0 . This result of this is shown in figure 3(b)

Naively one could expect that the cutoff scale Q_0 is larger for more massive hadrons. The most naive guess would be a linear dependence of the form $Q_0 \approx \Lambda_{\text{eff}} + m_{\text{hadron}}$. Figure 3(b) shows that this is not correct and that at least a separate treatment of mesons and baryons is necessary. A linear dependence of Q_0 on e.g. the hadron mass per constituent quark already describes the data slightly better, as can be seen in figure 3(c).




Figure 4: The experimental values of ξ_p^* versus $\sqrt{s}$ for various identified hadrons. The curves are from the combined fit of Λ_{eff} and $F(\text{hadron})$ to the predicted dependence (equations (1) and (8)). From top to bottom, the curves correspond to pions ($\pi^\pm$, π^0), all charged particles, kaons ($K^\pm$, K^0), protons and Λ baryons.

5 Dependence of ξ_p^* on $\sqrt{s}$

A more quantitative comparison is possible for the scaling behaviour of ξ_p^* , as parametrised by the QCD scale Λ_{eff} . The dependence on $\sqrt{s} = 2E_{\text{beam}}$ of the experimental values of ξ_p^* from table 1 is presented in the figure 4.

The experimental values can be fitted easily to the dependence given by equation (1) with $F = 0$, or to a linear dependence on Y . The results for different types of hadrons are given in table 2. Both functions are acceptable and relation (1) gives approximately the correct slope, that should be equal for all hadrons.

Before going on, one should note that in this case the quality of the fits can be considered as an internal check of the estimated errors. The 14 parameter

particle	a_h	b_h	$\chi^2/\text{n.d.f.}$	Q'_0	$\chi^2/\text{n.d.f.}$	points
charged	0.75 ± 0.11	0.644 ± 0.029	1.24	0.211 ± 0.007	1.03	6
$\pi^\pm$	0.75 ± 0.08	0.687 ± 0.026	1.27	0.181 ± 0.005	1.61	8
π^0	0.72 ± 0.21	0.728 ± 0.053	0.52	0.146 ± 0.008	0.95	6
$\pi^\pm, \pi^0$	0.69 ± 0.07	0.715 ± 0.024	1.33	0.175 ± 0.004	2.15	14
$K^\pm$	0.29 ± 0.11	0.577 ± 0.035	2.05	0.589 ± 0.027	2.24	7
K^0	0.46 ± 0.12	0.529 ± 0.035	2.69	0.612 ± 0.023	5.91	9
$K^\pm, K^0$	0.42 ± 0.07	0.540 ± 0.022	2.14	0.603 ± 0.018	4.02	16
p	0.33 ± 0.15	0.562 ± 0.046	0.33	0.621 ± 0.032	1.07	7
Λ	0.03 ± 0.07	0.610 ± 0.019	0.34	0.770 ± 0.025	1.37	7
p, Λ	0.12 ± 0.07	0.596 ± 0.018	2.08	0.725 ± 0.020	2.08	14
total $\chi^2/\text{n.d.f.}$			1.35		1.94	

Table 2: Result of a fit of the experimental values of ξ_p^* in table 1 to a linear function $\xi_p^* = a_h + b_h \cdot \log(E_{cms}/\text{GeV})$ for various identified hadrons. The scale Q'_0 is the result of a second fit to the function $\xi_p^* = \frac{1}{2}Y + \sqrt{CY} - C$, where $Y \equiv \log(E_{beam}/Q'_0)$ and $C = 0.2915$. The fits improve slightly when data at the Υ are excluded. To compare the overall quality of the fits to the two functions, a total $\chi^2/\text{n.d.f.}$ is given, based on the fits for $\pi^\pm, \pi^0, K^\pm, K^0, p$, and Λ .

fit to separate linear functions of $\log E$ for each hadron leads to a total $\chi^2/\text{n.d.f.} = 1.35$. One could conclude that some of the (systematic) errors have been slightly underestimated. It can also be noted that the K^0 data give the largest contribution to χ^2 .

In the fit to equation (1) it is not easy to disentangle Λ_{eff} from $F(Q_0)$, since they are strongly correlated. The simple way out was to constrain $\Lambda_{\text{eff}} = Q_0 = Q'_0$, so that $F(\lambda) = 0$. The fitted values of Q'_0 are not equal to the corresponding values of Q_0 shown in figure 3. The values of Q'_0 are quite near to the particle mass for mesons, but for the baryons Q'_0 is significantly smaller.

Figure 5 shows the dependence of Q'_0 on (a) the hadron mass and (b) the hadron mass per valence quark. Neither of these plots shows a clear linear dependence. It is possible that this is again due to F in equation (1).

An alternative to setting $F = 0$ is to perform a simultaneous fit of equation (1) to all the data from table 1, where Λ_{eff} is a universal scale, while F is allowed to have different values for hadrons, pions, kaons and the two baryons.

Such a six parameter fit was performed by minimising χ^2 using the MINUIT [24] program, with zero as starting values for F_h . There is a strong correlation in the fit of $\log \Lambda_{\text{eff}}$ and $\bar{F}$, the average of F_h . Therefore it is practical to fit $\bar{F}$ and differences $F_h - \bar{F}$. The fit has 44 degrees of freedom and leads to $\chi^2/44 = 1.98$, corresponding to a probability of $1.1 \cdot 10^{-4}$. This value of χ^2 can also be compared to the values of $\chi^2/\text{n.d.f.}$ in table

2, where separate fits for every hadron type were done. This comparison shows that one should probably not reject equation (1) on the basis of the low probability, because it cannot be excluded that the errors of ξ_p^* in table 1 have been underestimated by a factor of $\sqrt{\chi^2/\text{n.d.f.}} = 1.41$, on average. Figure 4 shows that the fitted functions are very near to straight lines and that they follow the data points rather well. Addition of an $\mathcal{O}(Y^{3/2})$ term as in equation (1) does not improve the fit significantly.

The resulting parameters and their external errors are given by:


$$\begin{aligned}
\Lambda_{\text{eff}} &= 0.07^{+0.09}_{-0.05} \text{ GeV} \\
\bar{F} &= -1.14^{+0.53}_{-0.70} \\
F_\pi - \bar{F} &= 0.54 \pm 0.02 \\
F_{\text{charged}} - \bar{F} &= 0.43 \pm 0.02 \\
F_K - \bar{F} &= -0.26 \pm 0.02 \\
F_p - \bar{F} &= -0.28 \pm 0.03 \\
F_\Lambda - \bar{F} &= -0.42 \pm 0.02
\end{aligned} \tag{8}$$

The correlation coefficient of Λ_{eff} and $\bar{F}$ is 100%, while the values of $F_h - \bar{F}$ are constrained to have a zero sum.


6 The theoretical meaning of Λ_{eff}

The determined value of Λ_{eff} from (8) is related to the (running) strong coupling constant by:

$$\frac{\alpha_s}{2\pi} = \frac{1}{b Y} = \frac{1}{b \log(E_{\text{beam}}/\Lambda_{\text{eff}})}, \tag{9}$$



(a) Q'_0 versus the hadron mass



(b) Q'_0 versus the hadron mass per quark

Figure 5: The values of Q'_0 from table from 2 versus (a) the hadron mass, and (b) the mass per valence quark.

giving at the Z^0 :

$$\alpha_s(\text{LEP}) = 0.108 \pm 0.018 \quad (10)$$

This value is compatible with the more accurate determinations done e.g. at LEP. This would suggest that LPHD + MLLA can really give a quantitative description of the low x_p data, based on perturbative QCD.

However, theoretically the QCD scale Λ_{eff} of MLLA is not well defined. This means that the value of Λ_{eff} as determined in (8) can not be compared to known values of e.g. $\Lambda_{\overline{\text{MS}}}$. A measurement

of Λ_{QCD} in a well defined subtraction scheme is equivalent to a measurement of $\alpha_s(E_{\text{beam}})$ at a well defined energy scale. To fix this energy scale one would need the $\mathcal{O}(Y^{3/2})$ -term in equation 1.

To see this [23], one can express Y in equation (1) in terms of the running coupling $\alpha_s(E_{\text{beam}})$. Using equation (9), the scale dependence of equation (1) translates to:

$$\frac{d\xi_p^*(E_{\text{beam}}, Q_0)}{d \log E_{\text{beam}}} = \frac{1}{2} + \frac{a}{8N_c} \sqrt{\frac{N_c \alpha_s}{2\pi}} + \mathcal{O}(\alpha_s^{3/2}) \quad (11)$$

The $\sqrt{\alpha_s}$ term is the next-to-leading (MLLA) correction to the leading term. The next-to-next-to-leading term $\mathcal{O}(\alpha_s)$ vanishes. An uncertainty of the energy scale would lead to e.g. $\alpha_s(2E)$ instead of $\alpha_s(E)$, inducing a correction in equation (11) of $\mathcal{O}(\alpha_s^{3/2}(E))$. This shows that the $\mathcal{O}(\alpha_s^{3/2}(E))$ needs to be known for the energy scale to be defined exactly. This is the reason for talking about the scale Λ_{eff} , in stead of e.g. $\Lambda_{\overline{\text{MS}}}$.

7 Conclusion

The data of LEP are giving a wealth of information about the fragmentation of quarks and gluons, as well as the mechanisms at play in the hadronisation. Although Monte Carlo models like JETSET and HERWIG can describe these data very accurately, it is questionable whether these models with all their free parameters lead to a better understanding of the physics behind the formation of hadrons in a jet, or only to a better parametrisation.

The ‘LPHD + MLLA’ approach [1–4] is an interesting attempt to understand some properties of multihadron production in jets in terms of perturbative QCD. This approach tries to take perturbative QCD as near as possible to the limits posed by confinement. It gives predictions for properties of momentum spectra that can be compared with experimental data, but concentrating on observables that can be described without too many unknown free parameters.

The experimental values of the parameter ξ_p^* for various hadrons and at various centre of mass energies have been determined from the published momentum spectra. Subsequently, an investigation was made of the dependence of ξ_p^* on both the centre of mass energy and the mass of the identified

hadrons, and this could be compared with the theoretical predictions.

The dependence of ξ_p^* on the centre of mass energy can be described adequately by the MLLA calculations. It is perhaps striking that the low x_p data, that are so intimately related to confinement, can not only be described by perturbative QCD, but can even be used to extract a consistent value of the strong coupling constant. The dependence of ξ_p^* on the mass and flavour of the identified hadron raises some questions that have not yet been answered in the context of LPHD + MLLA and the description of fragmentation by truncated parton cascades.

Acknowledgements

The author thanks NIKHEF-H for their hospitality while this work was finished and Bert Koene and Paul Kooijman for their critical comments. He is grateful for helpful comments and suggestions from Yuri L. Dokshitzer and Valery A. Khoze.

References

- [1] Yu.L. Dokshitzer, V.A. Khoze, A.H. Mueller and S.I. Troyan, ‘*Basics of Perturbative QCD*’, Editions Frontières, Gif-sur-Yvette, France, 1991
- [2] Yu.L. Dokshitzer, V.A. Khoze and S.I. Troyan, Journ. Mod. Phys. A 7 (1992) 1875.
- [3] Yu.L. Dokshitzer, V.A. Khoze and S.I. Troyan, Z. Phys. **C55** (1992) 107-114
- [4] Yu.L. Dokshitzer, V.S.Fadin and V.A.Khoze, Phys. Lett. **115B** (1982) 242; Z. Phys. **C15** (1982) 325; **C18** (1983) 37
Ya.I. Azimov, Yu.L.Dokshitzer, V.A.Khoze, and S.I.Troyan, Z. Phys. **C27**(1985) 65-72 ; **C31** (1986) 213-218
Yu.L. Dokshitzer, V.A. Khoze, C.P. Fong and B.R. Webber, Phys. Lett. **273B** (1991) 319-322
- [5] the OPAL collaboration, Phys. Lett. **247B** (1990) 617
- [6] the L3 collaboration, Phys. Lett. **259B** (1991) 199
- [7] E.R. Boudinov, P.V. Chliapnikov and V.A. Uvarov, Phys. Lett. **309B** (1993) 210
- [8] the DELPHI collaboration, Phys. Lett. **275B** (1992) 231
- [9] the OPAL collaboration, Phys. Lett. **264B** (1991) 467
- [10] the L3 collaboration, Phys. Lett. **286B** (1992) 403
- [11] the OPAL collaboration, Phys. Lett. **291B** (1992) 503
- [12] N.C. Brümmer, ‘*the Λ baryon as a probe of QCD at LEP*’, PhD thesis, NIKHEF-H Amsterdam, University of Leiden, the Netherlands (1994)
- [13] C.P. Fong and B. Webber, Phys. Lett. **229B** (1989) 289; Nucl. Phys. **B355** (1991) 54
- [14] A. Stuart and J.K. Ord, ‘*Kendall’s advanced theory of statistics*’, Vol.1, Griffin, 1987
- [15] the CLEO collaboration at CESR, Phys. Rev **D31** (1985) 2161; Phys. Rev. **D45** (1992) 752; Phys. Rev. Lett. **48** (1982) 1070; Phys. Rev. Lett. **58** (1987) 1814;
- [16] the ARGUS collaboration at DORIS, Z. Phys. **C39** (1988) 177; Z. Phys. **C42** (1989) 519; Z. Phys. **C44** (1989) 547; Z. Phys. **C46** (1990) 15; Z. Phys. **C58** (1993) 191
- [17] the TPC collaboration at PEP, Phys. Rev. **D52**(1984) 577; Phys. Rev. **D53**(1984) 2378; Phys. Rev. **D54**(1985) 274; Phys. Rev. Lett. **53** (1984) 2378; Phys. Rev. Lett. **54** (1985) 274; Phys. Rev. Lett. **61** (1988) 1263;
- [18] the HRS collaboration at PEP, Phys. Rev. **D35** (1987) 2639; Phys. Rev. **D45** (1992) 3949
- [19] the TASSO collaboration at PETRA, Phys. Lett. **94B** (1980) 444; Z. Phys. **C17** (1983) 5; Z. Phys. **C22** (1984) 307; Z. Phys. **C27** (1985) 27; Z. Phys. **C41** (1988) 359; Z. Phys. **C42** (1989) 189; Z. Phys. **C45** (1989) 209; Z. Phys. **C47** (1990) 187
- [20] the JADE collaboration at PETRA, Z. Phys. **C46** (1990) 1; **C28** (1985) 343
- [21] the TOPAZ collaboration, KEK annual report 1991, p. 54, 55
- [22] the DELPHI collaboration, Phys. Lett. **318B** (1993) 249.
- [23] private communications with Yuri L. Dokshitzer
- [24] F. James and M. Roos, CERN program library D506, the MINUIT - Function Minimisation and Error Analysis